arXiv:1512.07439v2 [math.RA] 1 Feb 2016

HOPF DENSE GALOIS EXTENSIONS WITH APPLICATIONS

JI-WEI HE, FRED VAN OYSTAEYEN AND YINHUO ZHANG

Abstract. Let H be a finite dimensional Hopf algebra, and let A be a left H-module algebra. Motivated by the study of the isolated singularities of A^H and the endomorphism ring $\mathrm{End}_{A^H}(A)$, we introduce the concept of Hopf dense Galois extensions in this paper. Hopf dense Galois extensions yield certain equivalences between the quotient categories over A and A^H. A special class of Hopf dense Galois extensions consits of the so-called densely group graded algebras, which are weaker versions of strongly graded algebras. A weaker version of Dade's Theorem holds for densely group graded algebras. As applications, we recover the classical equivalence of the noncommutative projective schemes over a noetherian $\mathbb{N}$-graded algebra A and its d-th Veroness subalgebra $A^{(d)}$ respectively. Hopf dense Galois extensions are also applied to the study of noncommuative graded isolated singularities.

Introduction

The study of Hopf algebra actions on Artin-Schelter regular algebras is always an important subject in the field of noncommutative algebraic geometry. Many interesting results are obtained in recent years (for instance, [Ue, MU, CKWZ1, CKWZ2, KKZ1, KKZ2, KKZ3, EW, WW] etc.). Let H be a Hopf algebra, and let A be a left H-module algebra. Assume A is also an Artin-Schelter regular algebra. In general, the invariant subalgebra A^H of A is not regular. Indeed, it was proved in [KKZ2] that the invariant subalgebra of a finite group action on a quantum polynomial algebra is regular if and only if the group is generated by quasi-reflections. If the Hopf algebra action satisfies some good conditions, say the homological determinant of H is trivial (for the terminology, see [JZ, KKZ1]), then the invariant subalgebra A^H is Artin-Schelter Gorenstein (cf. [KKZ1, KKZ3]). So, in this case, one asks about the properties of the singularities of A^H. Ueyama called a noetherian connected graded algebra A a *graded isolated singularity* if the quotient category $\mathrm{tail}\, A$ has finite global dimension (cf. [Ue]). He proved in [Ue] that a finite group action on an Artin-Schelter regular algebra of global dimension 2 with trivial homological determinant always yields a graded isolated singularity.

Let G be a finite group which acts homogeneously on an Artin-Schelter regular algebra A. If the invariant subalgebra A^G is a graded isolated singularity, then there is an equivalence of abelian categories $\mathrm{tail}\, A * G \cong \mathrm{tail}\, A^G$ (cf. [Ue, MU]), where $A * G$ is the skew group algebra of A and G. More generally, Mori and Ueyama call a finite group action on A *ample* if there is an equivalence $\mathrm{tail}\, A * G \cong \mathrm{tail}\, A^G$ (cf. [MU]). We found that the whole theory of graded isolated singularities and ample group actions on Artin-Schelter regular algebras fits into a more general theory, *Hopf dense Galois extensions*, which we will introduce in this paper. Our aim of this paper is to develop

2010 *Mathematics Subject Classification.* 16T05, 16W22.

Key words and phrases. Hopf dense Galois extension, densely graded algebra, quotient category.

a general theory on Hopf dense Galois extensions, and then in this framework to understand the finite dimensional Hopf-actions on Artin-Schelter regular algebras.

Let H be a finite dimensional Hopf algebra, and A a left H-module algebra. We may view A as a right H^*-comodule algebra with a coaction map $\rho : A \to A \otimes H^*$. Recall that the algebra extension A/A^H is called a *right H^*-Galois extension* if the map $\beta : A\otimes_{A^H} A \to A\otimes H^*, a\otimes b \mapsto \sum_{(b)} ab_{(0)}\otimes b_{(1)}$ is an isomorphism, where $\rho(b) = \sum_{(b)} b_{(0)} \otimes b_{(1)}$. When we study H-actions on infinite dimensional algebras, say Artin-Schelter regular algebras, the canonical map β is not often an isomorphism. So, we have to modify the definition of the classical Hopf Galois extension. We require that the map β is almost surjective, that is, the cokernel of β is finite dimensional. In this case, we call A/A^H a *right H^*-dense Galois extension* (for more details, see Section 1). It turns out that this is a efficient way to study the noncommutative projective schemes over the invariant subalgebra A^H, especially when A^H is a graded isolated singularity.

The paper is organized as follows. In Section 1, we introduce the definition of Hopf dense Galois extensions. In Section 2, we prove some general equivalences of abelian categories for Hopf dense Galois extensions. Let H be a finite dimensional semisimple Hopf algebra, A be a left H-module algebra. In the theory of the classical Hopf Galois extensions, the module ${}_{A\#H}A$ is a projective generator of the category of left $A\#H$-modules, and $\mathrm{End}(A_{A^H}) \cong A\#H$ as algebras (cf. [CFM, Theorem 1.9]). However, for the Hopf dense Galois extensions, these properties are not true in general. We need more restrictions on the algebra A. In Section 3, we show that if A satisfies some additional homological properties, then we have an isomorphism $\mathrm{End}(A_{A^H}) \cong A\#H$. The main results of Sections 2 and 3 are the following theorem (cf. Theorems 2.3 and 3.8). The notions in the theorem will be introduced in the corresponding sections.

Theorem. *Let H be a finite dimensional semisimple Hopf algebra, and let A be a left H-module algebra. Assume that A is also (on both sides) noetherian. Let $B = A\#H$, $R = A^H$ and t a nonzero integral of H. The following statements* (i) *to* (iv) *are equivalent.*

(i) *A/R is a right H^*-dense Galois extension.*
(ii) *For any finitely generated right B-module M, $T(M)$ is finite dimensional, where $T(M)$ the largest t-torsion submodule of M (cf. Section 2).*
(iii) *$B/(AtA)$ is finite dimensional.*
(iv) *$-\otimes_{\mathcal{B}} \mathcal{A} : \mathrm{QMod}\, B \longrightarrow \mathrm{QMod}\, R$ is an equivalence of abelian categories.*

If we assume further that $\mathrm{depth}\, A_A \geq 2$*, then the above statements implies that*

(v) *the natural map*

$$A\#H \longrightarrow \mathrm{End}(A_{A^H}), a\#h \mapsto [b \mapsto a(h\cdot b)]$$

is an isomorphism of algebras (cf. Section 3).

The statement (v) above follows from a more general result about equivalences of quotient categories (cf. Theorem 3.6). We remark that a part of the theorem above was obtained in [MU, Theorem 2.13] in the case that H is a finite group algebra and A is $\mathbb{N}$-graded.

A special class of Hopf dense Galois extensions are the so called *densely group graded algebras* which we will introduce in Section 4. It is a weaker version of strongly graded algebras (cf. [NV]). We have the following version of Dade's Theorem (cf. Theorem 4.7).

Theorem. *Let G be a finite group, and let $A = \oplus_{g\in G} A_g$ be a G-graded algebra. Assume that A is a noetherian algebra. Then the following are equivalent.*

(i) *A is a densely G-graded algebra.*
(ii) *For any finitely generated G-graded A-module $M = \oplus_{g\in G} M_g$, if M_e is finite dimensional, then M itself is finite dimensional.*
(iii) *The functor $(-)_e : \mathrm{QGrMod}_G A \longrightarrow \mathrm{QMod}\, A_e$ is an equivalence of abelian categories.*

For any $\mathbb{Z}$-graded algebra $A = \oplus_{i\in\mathbb{Z}} A_i$, we may view A as a $\mathbb{Z}_d$-graded algebra. Indeed, if we set $B_{\overline{i}} = \oplus_{r\in\mathbb{Z}} A_{rd+i}$ for all $\overline{i} \in \mathbb{Z}_d$ $(0 \leq i \leq d-1)$, then $B = \oplus_{\overline{i}\in\mathbb{Z}_d} B_{\overline{i}}$ is a $\mathbb{Z}_d$-graded algebra. Note that $B_{\overline{0}} = A^{(d)} = \oplus_{r\in\mathbb{Z}} A_{rd}$ is the dth Veroness subalgebra of A. As an applications of densely graded aglebras, in Section 5 we recover the classical result about the equivalence of the categories of the noncommutative projective schemes $\mathrm{tail}\, A^{(d)}$ and $\mathrm{tail}\, A$. Indeed, we have the following more general result (cf. Theorem 5.5).

Theorem. *Let $A = A_0 \oplus A_1 \oplus \cdots$ be a locally finite noetherian $\mathbb{N}$-graded algebra and let d be a fixed positive integer. The following are equivalent.*

(i) *There is an integer $p > 0$ such that, for all $n \geq p$ and $0 \leq s \leq d-1$,*

$$A_{nd} = \sum_{i+j=n-1} A_{id+s} A_{jd+d-s}.$$

(ii) *Given a finitely generated right graded A-module M, if $M^{(d)} = \oplus_{n\in\mathbb{Z}} M_{nd}$ is finite dimensional, then M itself is finite dimensional.*
(iii) *$(-)^{(d)} : \mathrm{Tail}\, A \longrightarrow \mathrm{Tail}\, A^{(d)}$ is an equivalence of abelian categories.*
(iv) *$(-)^{(d)} : \mathrm{tail}\, A \longrightarrow \mathrm{tail}\, A^{(d)}$ is an equivalence of abelian categories.*

Note that the condition (i) in the above theorem is relatively easy to verify. For example, if the graded algebra A is generated by elements in A_0 and A_1, then (i) is satisfied. Hence we recover the classical result (cf. Theorem 5.1). We remark that Mori also provided some equivalent conditions for the equivalence of $\mathrm{tail}\, A$ and $\mathrm{tail}\, A^{(d)}$ in [Mo, Theorem 3.5]. Note that it is assumed in [Mo] that A is a connected graded algebra satisfying further homological conditions. In our case, we only assume that A is a locally finite noetherian algebra, and we prove the result in the framework of Hopf dense Galois extension, which seems to be a new way to understand noncommutative projective schemes.

In Section 6, we provide another application of Hopf dense Galois extensions to graded isolated singularities. We prove that the invariant subalgebra A^H is a graded isolated singularity if A/A^H is a H^*-dense Galois extension and A has finite global dimension (cf. Corollary 6.2). Some further results about the endomorphism ring of A_{A^H} are obtained.

Throughout the paper, $\Bbbk$ is field of characteristic zero. All the algebras and modules are over $\Bbbk$, and the symbol $\otimes$ we means $\otimes_\Bbbk$. For the basic properties of classical Galois extensions, we refer to the references [CFM] and [Mon]. For the basic properties of quotient categories, we refer to the books [St] and [PP].

1. Hopf dense Galois extensions

Let H be a Hopf algebra with the coproduct Δ and antipode S. Let A be a right H-comodule algebra, that is, A is an algebra together with a right H-comodule action $\rho : A \to A \otimes H$, satisfying the compatibility condition?

$$\rho(ab) = \sum_{(b)(a)} a_{(0)}b_{(1)} \otimes a_{(1)}b_{(2)},$$

where we use Sweedler's sigma notation. The coinvariants of the H-coaction are defined to be $A^{coH} = \{a \in A | \rho(a) = a \otimes 1\}$, which is a subalgebra of A. If H is finite dimensional, the H^* is a Hopf algebra. The right H-coaction on A induces a left H^*-action by setting $f \cdot a = \sum_a f(a_{(1)})a_{(0)}$. Then A is a left H^*-algebra. The invariant subalgebra of H^*-action on A is defined to be $A^{H^*} = \{a \in A | f \cdot a = f(1)a\}$. One has $A^{coH} = A^{H^*}$ in case H is finite dimensional.

Recall from [KT] that the algebra extension A/A^{coH} is said to be *right H-Galois* if the map

$$\beta : A \otimes_{A^{coH}} A \to A \otimes H, a \otimes b \mapsto (a \otimes 1)\rho(b)$$

is surjective. In this case, A^{coH} shares many common properties with the smash product $A\#H^*$. Indeed, A^{coH} is Morita equivalent to $A\#H^*$ (cf. [CFM]) when H is semisimple. In many interesting examples in noncommutative algebraic geometry, the algebra extension A/A^{coH} may not be H-Galois, but still the coinvariant subalgebra A^{coH} shares some common properties with $A\#H^*$. It seems necessary to introduce a more general definition than the one of an H-Galois extension.

We first introduce some terminologies. Let B be an algebra, and M and N left B-modules. Let $f : M \to N$ be a left B-module morphism. We say that f is *almost surjective* if the image of f is cofinite, that is, $N/\operatorname{im}(f)$ is finite dimensional. We say that f is *almost injective* if $\ker f$ is finite dimensional. If f is both almost surjective and almost injective, we say that f is an *almost isomorphism*.

Definition 1.1. *Let H be a Hopf algebra, and let A be a right H-comodule algebra. We say that the algebra extension A/A^{coH} is* right H-dense Galois extension *if the map*

$$\beta : A \otimes_{A^{coH}} A \to A \otimes H, a \otimes b \mapsto (a \otimes 1)\rho(b)$$

is almost surjective.

Note that if A is finite dimensional, then any coaction of a finite dimensional Hopf algebra on A yields a H-dense Galois extension. Hence the definition is trivial for finite dimensional comodule algebras. When the comodule algebra A is infinite dimensional, there will be many interesting examples in noncommutative algebraic geometry. We give the following example, more examples will be given in following sections.

Example 1.2. *Let* $\sigma = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ *and let $G = \langle\sigma\rangle$ be the cyclic group of order 2. Let $A = \Bbbk[x, y]$ be the polynomial algebra. We view A as a $\mathbb{N}$-graded algebra. Then G acts homogeneously on A . Let $\Bbbk G$ be the group algebra, and let $H = \Bbbk G^*$. Then A is a right H-comodule algebra. One sees that $A^{coH} = \oplus_{n\geq 0} A_{2n}$. Now it is easy to see that $\dim((A \otimes H)/\operatorname{im}\beta) = 1$. Hence A/A^{coH} is a right H-dense Galois extension.*

Similar to the case of H-Galois extensions, we have the following property of H-dense Galois extensions.

Proposition 1.3. *Let H be a finite dimensional Hopf algebra, and let A be a left H-module algebra. Then A/A^H is a right H^*-dense Galois extension if and only if the map $[\ ,\ ] : A \otimes_{A^H} A \to A\#H, a \otimes b \mapsto (a\#t)(b\#1)$ is almost surjective, where t is a nonzero integral of H.*

Proof. Since H is finite dimensional, the map $\theta : H^* \to H$ defined by $\theta(f) = t \leftharpoonup f$ is a right H^*-module isomorphism. As showed in the proof of [CFM, Theorem 1.2], the map $[\ ,\ ] = (\mathrm{id} \otimes \theta) \circ \beta$. Hence $[\ ,\ ]$ is almost surjective if and only if β is almost surjective.

2. General equivalences of quotient categories

Let B be a noetherian algebra. We write $\operatorname{Mod} B$ for the category of right B-modules, and $\operatorname{mod} B$ for the full subcategory of $\operatorname{Mod} B$ consisting of finitely generated right B-modules. Let M_B be a right B-module. We call M a *torsion module* if for all $m \in M$, the submodule mB is finite dimensional. Note that every B-module admits a largest torsion submodule. We write $\operatorname{Tor} B$ for the full subcategory of $\operatorname{Mod} B$ consisting of all the torsion modules. Let $\operatorname{tor} B$ be the intersection of $\operatorname{mod} B$ and $\operatorname{Tor} B$.

Since B is noetherian, we see that $\operatorname{Tor} B$ (resp. $\operatorname{tor} B$) is a Serre subcategory of $\operatorname{Mod} B$ (resp. $\operatorname{mod} B$). Define

$$\operatorname{QMod} B := \frac{\operatorname{Mod} B}{\operatorname{Tor} B}, \qquad \operatorname{qmod} B := \frac{\operatorname{mod} B}{\operatorname{tor} B}.$$

Then both $\operatorname{QMod} B$ and $\operatorname{qmod} B$ are abelian categories.

Throughout the rest of this section, let H be a finite dimensional semisimple Hopf algebra, and let A be a left H-module algebra. Set $R = A^H$ to be the invariant subalgebra of A, and set $B := A\#H$. Assume that A is noetherian. Then both B and R are noetherian, moreover, A viewed as a left (or a right) R-module is finitely generated (cf. [Mon]).

We introduce another torsion class in the category of B-modules. Let $t \in H$ be the integral in H such that $\varepsilon(t) = 1$. Hence $t^2 = t$. The ideal of B generated by t is equal to $(A\#t)(A\#1)$, and hence is usually written by AtA. We call a right B-module M a *t-torsion module* if for every element $m \in M$ we have $m(A\#t)$ is finite dimensional. Let $\mathcal{T}$ be the full subcategory of $\operatorname{Mod} B$ consisting of all the t-torsion modules. By the definition, we see that $\operatorname{Tor} B$ is a full subcategory of $\mathcal{T}$. Let N be a right B-module. Write $T(N)$ for the sum of all the t-torsion submodules of N. Then $T(N)$ is the largest t-torsion submodule of N. Hence, we indeed obtain a functor

$$T : \operatorname{Mod} B \to \mathcal{T}.$$

Note that we have $Mt \in \operatorname{Mod} R$ for any $M \in \operatorname{Mod} B$.

Lemma 2.1. (i) *If $M \in \mathcal{T}$ is a finitely generated B-module, then $Mt = M(A\#t)$ is finite dimensional.*

(ii) *$M \in \mathcal{T}$ if and only if $Mt \in \operatorname{Tor} R$.*

(iii) *$\mathcal{T}$ is a localizing Serre subcategory of* $\operatorname{Mod} B$.

Proof. The first statement is clear.

(ii) Assume $M \in \mathcal{T}$. For $m \in M$, we have $(mt)R \subset (m(A\#H))t = m(A\#t)$. Hence the R-submodule of Mt generated by mt is finite dimensional. Therefore, $Mt \in \operatorname{Tor} R$. On the contrary, assume $Mt \in \operatorname{Tor} R$. By assumption, A is a noetherian algebra. Hence A, viewed as a right R-module, is finitely generated. So, $(A\#H)t = A\#t$ is a finitely generated R-module. Take an arbitrary element $m \in M$. We see that $m(A\#t)$ is finitely generated R-module. On the other hand, $m(A\#t) = m(A\#H)t \subset Mt$, impling that $m(A\#t)$ is a finitely generated module in $\operatorname{Tor} R$. Hence $m(A\#t)$ is finite dimensional. Therefore $M \in \mathcal{T}$.

(iii) One easily sees that $\mathcal{T}$ is closed under taking arbitrary direct sums. Let M be a B-module, and let K be a submodule module of M. Consider the exact sequence $0 \to K \hookrightarrow M \xrightarrow{g} N \to 0$. If $M \in \mathcal{T}$, then one easily sees both K and N are in $\mathcal{T}$. Assume K and N are in $\mathcal{T}$. Taking an element $m \in M$, we have that $g(m(A\#t)) = g(m)(A\#t)$ is finite dimensional. Since $A\#H$ is noetherian, $mB \cap K$ is a finitely generated B-module. Then we have that $m(A\#t) \cap K = m(A\#t) \cap Kt = (mB \cap K)t$ is finite dimensional. Now the exact sequence $0 \to m(A\#t) \cap K \to m(A\#t) \to g(m(A\#t)) \to 0$ implies that $m(A\#t)$ is finite dimensional. Hence M is in $\mathcal{T}$. □

Let $R = A^H$ be the invariant subalgebra of A, and let $\operatorname{Mod} R$ be the category of right R-module. Then ${}_RA_B$ and ${}_BA_R$ are bimodules. The functor $- \otimes_B A : \operatorname{Mod} B \longrightarrow \operatorname{Mod} R$ has a right adjoint functor $\operatorname{Hom}_R({}_BA_R, -)$. Since the functor $- \otimes_B A$ is isomorphic to $(-)^H$, one sees that there is a natural isomorphism $\operatorname{Hom}_R({}_BA_R, -) \circ (- \otimes_B A) \longrightarrow \mathrm{id}_{\operatorname{Mod} R}$.

Consider the quotient category $\operatorname{QMod} R = \frac{\operatorname{Mod} R}{\operatorname{Tor} R}$. Let $\pi : \operatorname{Mod} R \longrightarrow \operatorname{QMod} R$ be the natural projection functor. Since $\operatorname{Tor} R$ is a localizing subcategory, π has a right adjoint functor $\omega : \operatorname{QMod} R \longrightarrow \operatorname{Mod} R$. Thus we have the following functor of adjoint pairs:

$$\operatorname{Mod} B \underset{\operatorname{Hom}_R({}_BA_R, -)}{\overset{- \otimes_B A_R}{\rightleftarrows}} \operatorname{Mod} R \underset{\omega}{\overset{\pi}{\rightleftarrows}} \operatorname{QMod} R \ .$$

Let $F = \pi \circ (- \otimes_B A_R)$ and $G = \operatorname{Hom}_R({}_BA_R, -) \circ \omega$. Then (F, G) is a pair of adjoint functors. Since $(- \otimes_B A_R) \circ \operatorname{Hom}_R({}_BA_R, -)$ is isomorphic to $\mathrm{id}_{\operatorname{Mod} R}$ and $\pi \circ \omega$ is isomorphic to $\mathrm{id}_{\operatorname{QMod} R}$, we have that FG is isomorphic to $\mathrm{id}_{\operatorname{QMod} R}$. Since H is semisimple, ${}_BA$ is a projective module. Hence $- \otimes_B A_R$ is an exact functor. Therefore F is an exact functor. Then a classical result of torsion theory (cf. [PP, Theorem 7.11, Chapter 4]) shows that $\ker F$ is a localizing Serre subcategory of $\operatorname{Mod} B$ and F induces an isomorphism of abelian categories

$$\overline{F} : \frac{\operatorname{Mod} B}{\ker F} \cong \operatorname{QMod} R. \tag{1}$$

Let us check the objects in $\ker F$. For $M \in \operatorname{Mod} B$, we have $F(M) = \pi(M \otimes_B A) = \pi(Mt)$. Hence $F(M) = 0$ if and only if $Mt \in \operatorname{Tor} R$. By Lemma 4.6, $M \in \ker F$ if and only if $M \in \mathcal{T}$.

In summarizing, we have the following result, which may be viewed as a generalization of [VZ, Theorem 2.4], and also can be compared to [Ga, Theorem 4.6].

Theorem 2.2. *Let H be a finite dimensional semisimple Hopf algebra, and let A be a noetherian left H-module algebra. Then we have an equivalence of abelian categories*

$$\operatorname{Mod} B/\mathcal{T} \cong \operatorname{QMod} R.$$

Since $\operatorname{Tor} B$ is a full subcategory of $\mathcal{T}$, it is indeed a Serre subcategory of $\mathcal{T}$. Hence we have a quotient category $\mathcal{T}/\operatorname{Tor} B$. Moreover, $\mathcal{T}/\operatorname{Tor} B$ is a full subcategory of $\operatorname{QMod} B$. Since $-\otimes_B A$ is an exact functor, it induces an exact functor

$$-\otimes_{\mathcal{B}}\mathcal{A} : \operatorname{QMod} B \longrightarrow \operatorname{QMod} R. \tag{2}$$

The exact functor F defined above induces an exact sequence of abelian categories

$$0 \longrightarrow \mathcal{T}/\operatorname{Tor} B \longrightarrow \operatorname{QMod} B \xrightarrow{-\otimes_{\mathcal{B}}\mathcal{A}} \operatorname{QMod} R \longrightarrow 0. \tag{3}$$

Theorem 2.3. *Let A and H be as in Theorem 2.2. Let $B = A\#H$ and $R = A^H$. The following are equivalent.*

(i) *A/R is a right H^*-dense Galois extension;*
(ii) *For any finitely generated right B-module M, $T(M)$ is finite dimensional;*
(iii) *$B/(AtA)$ is finite dimensional;*
(iv) *$-\otimes_{\mathcal{B}}\mathcal{A} : \operatorname{QMod} B \longrightarrow \operatorname{QMod} R$ is an equivalence of abelian categories.*

Proof. The equivalence of (i) and (iii) is Proposition 1.3. That (ii) implies (iii) is obvious since $B/(AtA)$ is a t-torsion B-module.

(iii) $\Longrightarrow$ (ii). Let M_B be a finitely generated module. Since A is (and hence B is) noetherian, $T(M)$ is finitely generated. By Lemma 4.6, $T(M)t$ is finite dimensional R-module. Hence $T(M)t\otimes_R A$ is finite dimensional since A is finitely generated as a left R-module. Since the right multiplication map $T(M)t\otimes_R A \longrightarrow T(M)(AtA) = T(M)tA$ is surjective, it follows that $T(M)tA$ is finite dimensional. The right B-module $T(M)/(T(M)tA)$ is finitely generated, which also can be viewed as a finitely generated $B/(AtA)$-module. By the hypothesis (iii), $B/(AtA)$ is finite dimensional, hence $T(M)/(T(M)tA)$ is finite dimensional. Together with the property that $T(M)tA$ is finite dimensional, we obtain that $T(M)$ is finite dimensional.

(ii) $\Longrightarrow$ (iv). Take an object $M \in \mathcal{T}$. For an arbitrary element $m \in M$, mB is an object in $\mathcal{T}$. By the condition (ii), $mB = T(mB)$ is finite dimensional. Hence $M \in \operatorname{Tor} B$. Therefore $\operatorname{Tor} B = \mathcal{T}$. From the exact sequence (3), we obtain that $-\otimes_B A : \operatorname{QMod} B \longrightarrow \operatorname{QMod} R$ is an equivalence.

(iv) $\Longrightarrow$ (ii). The condition (iv) implies $\operatorname{Tor} B = \mathcal{T}$. If M be a finitely generated B-module, so is $T(M)$ since B is noetherian. Then $T(M) \in \mathcal{T} = \operatorname{Tor} B$ is finite dimensional. □

3. Endomorphism rings

Let H be a finitely dimensional semisimple Hopf algebra, and A be a left H-module algebra. Recall from [CFM, Theorem 1.2] that if A is a right H^*-Galois extension, then there is an isomorphism $\operatorname{End}_{A^H}(A) \cong A\#H$. In the Hopf dense Galois case, this property is not true in general. However, we will see that a similar property holds if we put more restrictions on A.

The discussions in this section fit in a more general setting. Let R be a noetherian algebra. Let $\operatorname{Mod} R$ be the category of right R-modules, and $\operatorname{Tor} R$ be the subcategory of torsion modules. Consider the torsion functor $\tau : \operatorname{Mod} R \longrightarrow \operatorname{Tor} R$ sending each module $M \in \operatorname{Mod} R$ to its largest torsion submodule. τ is a left exact functor. We write $R^i\tau$ for the ith right derived functor of τ.

Let M_R be a finitely generated R-module. We define the *depth* of M to be

$$\operatorname{depth}(M) = \min\{i | R^i\tau(M) \neq 0\}.$$

Lemma 3.1. *Let R be a noetherian algebra, and M_R a finitely generated R-module. Then the following are equivalent.*

(i) $\operatorname{depth}(M) \geq d$.
(ii) $\operatorname{Ext}^i_R(S, M) = 0$ *for all* $i < d$ *and all finite dimensional simple module* S_R.
(iii) $\operatorname{Ext}^i_R(K, M) = 0$ *for all* $i < d$ *and all finite dimensional module* K_R.

Proof. (i) $\Longleftrightarrow$ (ii). Take a minimal injective resolution of M as: $0 \longrightarrow M \longrightarrow I^0 \longrightarrow I^1 \longrightarrow \cdots \longrightarrow I^i \longrightarrow \cdots$. Assume $\operatorname{depth}(M) \geq d$. We claim that I^i is torsion free for all $i < d$. If $d = 0$, nothing needs to be proved. We assume $d > 0$. Since $R^0\tau(M) = \tau(M)$, we see M is torsion free. Hence the injective enveloping I^0 of M is also torsion free. Let i be the smallest number such that I^i is not torsion free. Then $Soc(I^i) \cap \tau(M) \neq 0$. Since the injective resolution is assumed to be minimal, we see $R^i\tau(M) \neq 0$, which contradicts to the assumption that $\operatorname{depth}(M) \geq d$. The claim follows. By the minimality of the injective resolution again, for any finite dimensional simple module S_R, we have $\operatorname{Ext}^i_R(S, M) = \operatorname{Hom}_R(S, I^i) = 0$ for all $i < d$.

The foregoing also holds that if we assume that $\operatorname{Ext}^i_R(S, M) \neq 0$ for all $i < d$ and all finite dimensional simple module S_R. Then I^i is torsion free for all $i < d$, which in turn implies $\operatorname{depth}(M) \geq d$.

(ii) $\Longleftrightarrow$ (iii). It suffices to show that (ii) implies (iii). Since any finite dimensional module K_R contains a simple submodule, the result follows from an easy induction on the dimension of K. □

Given noetherian algebras R and B and a B-R-bimodule ${}_BM_R$, we are interested in when M defines an equivalence between the quotient categories $\operatorname{QMod} R$ and $\operatorname{QMod} B$. We need some terminology.

Let ${}_BP$ be a B-module. We say that ${}_BP$ is *almost flat* if given finitely generated module K_B and N_B and an injective morphism $f : K_B \to N_B$, the kernel of the map $f \otimes_B P : K \otimes_B P \to N \otimes_B P$ is finite dimensional. We say that ${}_BP$ is *faithful* if for any finitely generated module N_B, the finite dimensionality of $N \otimes_B M$ implies that N_B itself is finite dimensional. We say ${}_BP$ is a *weaker generator* of the category of all the left B-modules, if there is a finite index set I and an almost surjective B-module morphism $\oplus_{i\in I}P_i \longrightarrow B/\tau(B)$, where P_i is a cofinite submodule of ${}_BP$ for all $i \in I$.

Lemma 3.2. *Let B be a noetherian algebra, and let ${}_BP$ be a finitely generated B-module. Then ${}_BP$ is almost flat if and only if for any almost injective B-module morphism $f : K_B \to N_B$ with K_B and N_B finitely generated, the kernel of the morphism $f \otimes_B P$ is finite dimensional.*

Proof. Assume that ${}_BP$ is almost flat. Let $f : K_B \to N_B$ be an almost injective morphism. We have an injective morphism $\overline{f} : K/\ker f \longrightarrow N$ induced by f. Then $\ker(\overline{f} \otimes_B P)$ is finite dimensional. Let $p : K \to K/\ker f$ be the natural projection map. Then the morphism $f \otimes_B P$ is the composition $K \otimes_B P \stackrel{p\otimes_B P}{\longrightarrow} K/\ker f \otimes_B P \stackrel{\overline{f}\otimes_B P}{\longrightarrow} N \otimes_B P$. We have an exact sequence of B-modules $0 \longrightarrow (\ker f) \otimes_B P \longrightarrow \ker(f \otimes_B P) \longrightarrow \ker(\overline{f} \otimes_B P) \longrightarrow 0$. Since f is almost injective, $\ker f$ is finite dimensional. Since ${}_BP$ is finitely generated, $(\ker f) \otimes_B P$ is finite dimensional. Hence $\ker(f \otimes_B P)$ is finite dimensional. □

Lemma 3.3. *Let R and B be noetherian algebras, and let ${}_BM_R$ be a B-R-bimodule. Assume that M is finitely generated on both sides, and that ${}_BM$ is a faithful and almost flat B-module. For a module N_B, $N \otimes_B M$ is a torsion R-module if and only if N_B is a torsion B-module.*

Proof. Assume that $N \otimes_B M$ is a torsion R-module. Let K_B be a finitely generated submodule of N_B, and let $\iota : K \to N$ be the inclusion map. Since ${}_BM$ is almost flat, $\ker(\iota \otimes_B M)$ is finite dimensional by Lemma 3.2. Since $N \otimes_B M$ is a torsion R-module, the image of $\iota \otimes_B M$ is a torsion R-module. Hence $K \otimes_B M$ is a torsion R-module. Since M is finitely generated as a right R-module, $K \otimes_B M$ is a finite generated R-module. Hence $K \otimes_B M$ is finite dimensional. By the assumption that ${}_BM$ is faithful, we obtain that K_B is finite dimensional. Therefore, N_B is a torsion module.

Conversely, assume that N_B is a torsion B-module. Note that $N = \varinjlim K$ where K runs over all the finite dimensional submodules of N_B. Since direct limits commute with left adjoint functor, we have $N \otimes_B M = (\varinjlim K) \otimes_B M \cong \varinjlim(K \otimes_B M)$. Since ${}_BM$ is finitely generated, $K \otimes_B M$ is finite dimensional for all finite dimension submodule K_B of N_B. Hence we obtain that $\varinjlim(K \otimes_B M)$ is a torsion R-module. □

Example 3.4. (i) Let B a noetherian algebra. Let ${}_BQ$ be a finitely generated projective B-module. If ${}_BP$ be a submodule of ${}_BQ$ such that Q/P is finite dimensional, then ${}_BP$ is an almost flat module. Indeed, let $f : X_B \to Y_B$ be an injective morphism. From the exact sequence $0 \longrightarrow P \xrightarrow{\iota} Q \xrightarrow{p} Q/P \longrightarrow 0$ then we have the following commutative diagram:

$$\begin{array}{ccccccc} Y \otimes_B P & \xrightarrow{Y\otimes_B\iota} & Y \otimes_B Q & \xrightarrow{Y\otimes_B p} & Y \otimes_B (Q/P) & \longrightarrow & 0 \\ \uparrow{\scriptstyle f\otimes_B P} & & \uparrow{\scriptstyle f\otimes_B Q} & & \uparrow{\scriptstyle f\otimes_B (Q/P)} & & \\ X \otimes_B P & \xrightarrow{X\otimes_B\iota} & X \otimes_B Q & \xrightarrow{X\otimes_B p} & X \otimes_B (Q/P) & \longrightarrow & 0. \end{array}$$

Hence we have $(f \otimes_B Q) \circ (X \otimes_B \iota) = (Y \otimes_B \iota) \circ (f \otimes_B P)$. Since Q is projective, $f \otimes_B Q$ is injective. Hence $\ker[(f \otimes_B Q) \circ (X \otimes_B \iota)] = \ker(X \otimes_B \iota)$. Since Q/P is finite dimensional and X_B is finitely generated, we have that $\mathrm{Tor}_1^B(X, Q/P)$ is finite dimensional. Then $\ker(X \otimes_B \iota)$ is finite dimensional. Therefore $\ker[(Y \otimes_B \iota) \circ (f \otimes_B P)]$ is finite dimensional, which implies that $\ker(f \otimes_B P)$ is finite dimensional.

(ii) Let B be a noetherian algebra. Let J be a left ideal of B. If B/J is finite dimensional, then J, as a left B-module, is faithful and almost flat. Indeed, J is almost flat by (i). Now assume X_B be a finitely generated B-module and $X \otimes_B J$ is finite dimensional. The exact sequence $X \otimes_B J \longrightarrow X \longrightarrow X \otimes_B (B/J) \longrightarrow 0$ implies that X is finite dimensional. Hence J is faithful.

Lemma 3.5. *Let R and B be noetherian algebras, and let ${}_BM_R$ be a B-R-bimodule which is finitely generated on both sides. Assume $g : X_B \to Y_B$ is an injective B-module morphism where X_B and Y_B are arbitrary B-modules. If M is almost flat as a left B-module, then the kernel of the morphism $g \otimes_B M$ is a torsion R-module.*

Proof. Note that $Y = \varinjlim Y'$ where the limit runs over all the finitely generated submodules Y' respectively. For any finitely generated submodule $Y' \subseteq Y$, set $X' = g^{-1}(Y' \cap g(X))$. Then we see $X = \varinjlim X'$. By Lemma 3.2, the kernel of the map $g|_{X'} : X' \otimes M \longrightarrow Y' \otimes_B M$ is finite dimensional, hence it is a torsion module. Let $K' = \ker(g|_{X'})$. For every finitely generated submodule Y' of Y,

we have an exact sequence $0 \longrightarrow K' \longrightarrow X' \otimes M \longrightarrow Y' \otimes_B M$. Taking direct limit on these exact sequence and using the property that direct limits are exact and commute with the tensor functor, we obtain that the kernel of $g \otimes_B M$ is a torsion R-module. □

Let ${}_BM_R$ be a B-R-bimodule which is finitely generated on both sides. Then we have functors $- \otimes_B M : \operatorname{Mod} B \longrightarrow \operatorname{Mod} R$, and $- \otimes_B M : \operatorname{mod} B \longrightarrow \operatorname{mod} R$. As the tensor functor $- \otimes_B M$ sends torsion B-modules to torsion R-modules and almost isomorphisms to almost isomorphisms, it induces functors $- \otimes_{\mathcal{B}} \mathcal{M} : \operatorname{QMod} B \longrightarrow \operatorname{QMod} R$ and $- \otimes_{\mathcal{B}} \mathcal{M} : \operatorname{qmod} B \longrightarrow \operatorname{qmod} R$, such that the following diagrams commute

$$\begin{array}{ccc} \operatorname{Mod} B & \xrightarrow{-\otimes_B M} & \operatorname{Mod} R \\ \downarrow{\scriptstyle \pi} & & \downarrow{\scriptstyle \pi} \\ \operatorname{QMod} B & \xrightarrow{-\otimes_{\mathcal{B}} \mathcal{M}} & \operatorname{QMod} R \end{array} \qquad \begin{array}{ccc} \operatorname{mod} B & \xrightarrow{-\otimes_B M} & \operatorname{mod} R \\ \downarrow{\scriptstyle \pi} & & \downarrow{\scriptstyle \pi} \\ \operatorname{qmod} B & \xrightarrow{-\otimes_{\mathcal{B}} \mathcal{M}} & \operatorname{qmod} R. \end{array}$$

Let M' be another B-R-bimodule. If there is a B-R-bimodule morphism $f : M \to M'$ which is an almost isomorphism, then it is not hard to see that f induces a natural isomorphism $- \otimes_{\mathcal{B}} \mathcal{M} \longrightarrow - \otimes_{\mathcal{B}} \mathcal{M}'$.

We have the following Morita type theorem for quotient categories.

Theorem 3.6. *Let R and B be noetherian algebras. Let ${}_BM_R$ be a bimodule which is finitely generated on both sides. Assume* $\operatorname{depth} M_R \geq 2$ *and* $\operatorname{depth} B_B \geq 2$. *Then the following are equivalent.*

(i) *The functor $- \otimes_{\mathcal{B}} \mathcal{M} : \operatorname{qmod} B \longrightarrow \operatorname{qmod} R$ is an equivalence of abelian categories.*
(ii) *M_R is a weaker generator, the natural map $B \to \operatorname{End}(M_R)$ is an isomorphism of algebras, and the left B-module ${}_BM$ is faithfully almost flat.*
(iii) *The functor $- \otimes_{\mathcal{B}} \mathcal{M} : \operatorname{QMod} B \longrightarrow \operatorname{QMod} R$ is an equivalence of abelian categories.*

Proof. (i) $\Longrightarrow$ (ii). Set $F = - \otimes_{\mathcal{B}} \mathcal{M}$. Then $F(\pi(B_B)) = \pi(M_R)$. Let I be a right ideal of B such that B/I is finite dimensional. Since $\operatorname{depth} B_B \geq 2$, we see $\operatorname{Hom}_B(B, B) \cong \operatorname{Hom}_B(I, B)$. Hence the projection functor $\pi : \operatorname{mod} B \longrightarrow \operatorname{qmod} B$ induces an isomorphism of algebras $\operatorname{End}(B_B) \xrightarrow{\pi} \operatorname{End}_{\operatorname{qmod} B}(\pi(B_B))$. Similar, since $\operatorname{depth} M_R \geq 2$, we have an isomorphism of algebra $\operatorname{End}(M_R) \xrightarrow{\pi} \operatorname{End}_{\operatorname{qmod} R}(\pi(M_R))$. By the commutative diagram before the theorem and the assumption that F is an equivalence of abelian categories, the natural map $B \to \operatorname{End}(M_R)$ is an isomorphism.

Assume $\pi(R) = F(\pi(N))$ for some $N \in \operatorname{mod} B$. Since N_B is finitely generated, there is a finite set Λ and an epimorphism $g : \oplus_{i \in \Lambda} B_i \longrightarrow N$, where $B_i \cong B$ for all $i \in \Lambda$. By the commutative diagrams before the theorem, we obtain an epimorphism $\varrho : \pi(\oplus_{i \in \Lambda} B_i \otimes_B M) \longrightarrow \pi(R)$ in $\operatorname{qmod} R$. Note that there is a cofinite submodule P of $\oplus_{i \in \Lambda} B_i \otimes_B M$ and a right R-module morphism $h : P \to R/\tau(R)$ such that $\pi(h) = \varrho$. Since ϱ is epimorphic, h is almost surjective. Write $M_i = B_i \otimes_B M \cong M$, and let p_i be the projection map $\oplus_{i \in \Lambda} M_i \to M_i$. Set $P_i = \operatorname{im}(p_i)$ for all $i \in \Lambda$. Since P is a cofinite submodule of $\oplus_{i \in \Lambda} M_i$, P_i is a cofinite submodule of M_i. Now we see the induced map $\oplus_{i \in \Lambda} P_i \longrightarrow R/\tau(R)$ is almost surjective. Hence M_R is a weaker generator.

For $K_B \in \operatorname{mod} B$, assume $K \otimes_B M$ is finite dimensional. Then we see $F(\pi(K)) \cong \pi(K \otimes_B M) = 0$. Since F is an equivalence, $\pi(K) = 0$. So, K_B is a torsion B-module, and hence K is finite dimensional. Therefore, ${}_BM$ is faithful. Let $g : X \to Y$ be an injective morphism in $\operatorname{mod} B$. From the commutative diagram before the theorem again, we see that $\pi(g \otimes_B M) : \pi(X \otimes_B M) \longrightarrow$

$\pi(Y \otimes_B M)$ is an injective morphism in $\operatorname{qmod} R$. Then $\ker(g \otimes_B M)$ is a torsion R-module, and hence $\ker(g \otimes_B M)$ is finite dimensional. Therefore ${}_BM$ is an almost flat module.

(ii) $\Longrightarrow$ (iii). Since M_R is a weaker generator in $\operatorname{Mod} R$, one sees that $\pi(M)$ is a generator in $\operatorname{QMod} R$. By Popescu-Gabriel's Theorem (cf. [St, Theorem X.4.1]), the functor $\operatorname{Hom}_{\operatorname{QMod} R}(\pi(M), -) : \operatorname{QMod} R \longrightarrow \operatorname{Mod} B$ is fully faithful, and has a left adjoint functor. Since π has a right adjoint functor $\omega : \operatorname{QMod} R \longrightarrow \operatorname{Mod} R$, one sees there is a natural isomorphism $\operatorname{Hom}_{\operatorname{QMod} R}(\pi(M), -) \cong \operatorname{Hom}_R(M, -) \circ \omega$. Hence the left adjunction of $\operatorname{Hom}_{\operatorname{QMod} R}(\pi(M), -)$ is $\pi \circ (- \otimes_B M) : \operatorname{Mod} B \longrightarrow \operatorname{QMod} R$. Write T for $\pi \circ (- \otimes_B M)$. Since ${}_BM$ is almost flat, T induces an equivalence of abelian categories $\operatorname{Mod} B/\ker T \longrightarrow \operatorname{QMod} R$, where $\ker T$ is the full subcategories of $\operatorname{Mod} B$ consists of objects K such that $T(K) = 0$. Since ${}_BM$ is faithful, we see $\ker T = \operatorname{Tor} B$ by Lemma 3.3. Hence $\operatorname{Mod} B/\ker T = \operatorname{QMod} B$. Therefore the functor $- \otimes_B \mathcal{M}$ induced by T is an equivalence.

(iii) $\Longrightarrow$ (i). This is clear since $\operatorname{qmod} B$ (resp. $\operatorname{qmod} R$) is a full subcategory of $\operatorname{QMod} B$ (resp. $\operatorname{QMod} R$) consisting of noetherian objects. □

Let us go back to the Hopf dense Galois extensions. Let H be a finite dimensional semisimple Hopf algebra, and let A be a left H-module algebra. We assume further that A is a noetherian algebra. As before, set $B := A\#H$ and $R := A^H$.

Lemma 3.7. *Let A and H be as above. If* $\operatorname{depth} A_A \geq 2$, *then* $\operatorname{depth} B_B \geq 2$ *and* $\operatorname{depth} A_R \geq 2$.

Proof. Let K_B be a finite dimensional B-module. We view K and B as right A-modules, then $\operatorname{Ext}^i_A(K, B)$ is a right H-module for all $i \geq 0$. By [HVZ, Corollary 2.9], we have $\operatorname{Ext}^i_B(K, B) = \operatorname{Ext}^i_A(K, B)^H$ for all $i \geq 0$, where $\operatorname{Ext}^i_A(K, B)^H$ is the subspace of invariants of H-actions. Since B is free A-module and $\operatorname{depth} A_A \geq 2$, we have $\operatorname{Ext}^i_A(K, B) = 0$ for all $i < 2$ by Lemma 3.1. Therefore $\operatorname{Ext}^i_B(K, B) = 0$ for all $i < 2$. Hence by Lemma 3.1 again $\operatorname{depth} B_B \geq 2$.

Let K_R be a finite dimensional module. We have the following isomorphisms $\operatorname{Hom}_B(K \otimes_R A_B, B) \cong \operatorname{Hom}_R(K, \operatorname{Hom}_B({}_RA_B, B)) \cong \operatorname{Hom}_R(K, {}_BA_R)$. These isomorphisms induce the following spectral sequence

$$E_2^{pq} = \operatorname{Ext}^p_B(\operatorname{Tor}^R_q(K, {}_RA_B), B) \Longrightarrow \operatorname{Ext}^{p+q}_R(K, {}_BA_R).$$

Since K_R is finite dimensional and ${}_RA$ is a finitely generated left R-module, $\operatorname{Tor}^R_q(K, {}_RA_B)$ is finite dimensional for all $q \geq 0$. Since $\operatorname{depth} B_B \geq 2$, we have $\operatorname{Ext}^p_B(\operatorname{Tor}^R_q(K, {}_RA_B), B) = 0$ for $p = 0, 1$ by Lemma 3.1. Hence $\operatorname{Ext}^i_R(K, {}_BA_R) = 0$ for $i = 0, 1$. By Lemma 3.1 again, $\operatorname{depth}(A_R) \geq 2$. □

As a special case of Theorem 3.6, we obtain the following result.

Theorem 3.8. *Let H be a finite dimensional semisimple Hopf algebra, and let A be a left H-module algebra such that A/A^H is right dense Galois. If A is noetherian and* $\operatorname{depth} A_A \geq 2$, *then the natural map*

$$A\#H \longrightarrow \operatorname{End}(A_{A^H}), a\#h \mapsto [b \mapsto a(h \cdot b)]$$

is an isomorphism of algebras.

Proof. This a direct consequence of Lemma 3.7 and Theorem 3.6. □

4. Densely graded algebras

Let G be a group, and let $A = \oplus_{g\in G} A_g$ be a G-graded algebra. Recall that A is called a *strongly graded algebra* if $A_gA_h = A_{gh}$ for all $g, h \in G$. We introduce a weaker version of strongly graded algebra, which is a special class of dense Galois extensions.

Definition 4.1. *Let G be a group, and let $A = \oplus_{g\in G} A_g$ be a G-graded algebra. We call A a* densely graded algebra *if* (i) *there are only finitely many elements $g \in G$ such that $A_gA_{g^{-1}} \neq A_e$, and* (ii) *$A_1/A_gA_{g^{-1}}$ is finite dimensional for all $g \in G$.*

Example 4.2. (i) If G is a finite group, then any finite dimensional G-graded algebra is densely graded.

(ii) Let $A = \Bbbk[x, y]/(x^2 - y^2)$, let $G = \mathbb{Z}_2$. We view the polynomial algebra $\Bbbk[x, y]$ as a G-graded algebra by setting the degree of x to be 0 and the degree of y to be 1. It follows that the element $x^2 - y^2$ is a homogeneous element of degree 0, which makes A as a G-graded algebra. We have $A_0 = \Bbbk[x]$ and $A_1 = y\Bbbk[x]$. So, $A_0^2 = A_0$, $A_1A_0 = A_0A_1 = A_1$ and A_0/A_1A_2 has dimension 2. Hence A is a densely graded G-graded algebra.

(iii) Let A be an algebra and let M be an A-A-bimodule. Assume $\varphi : M \otimes_A M \longrightarrow A$ is an A-A-bimodule morphism such that $A/\varphi(M \otimes_A M)$ is finite dimensional, and such that $m\varphi(n \otimes l) = \varphi(m \otimes n)l$. Set $B = (A \oplus M)$ and define multiplication on B as follows:

$$(a, m)(b, n) = (ab + \varphi(m \otimes n), an + mb), \text{ for all } a, b \in A \text{ and } n, m \in M.$$

Set $B_0 = A$ and $B_1 = M$. Then B is a densely $\mathbb{Z}_2$-graded algebra. We call B a *densely semitrivial extension* of A by M (compared to [NV, Example 1.3.3]).

The following observation is easy to check.

Lemma 4.3. *Let $A = \oplus_{g\in G} A_g$ be a G graded algebra. If $A_gA_h \neq A_{gh}$, then $A_gA_{g^{-1}} \neq A_e$ and $A_{h^{-1}}A_h \neq A_e$.*

We have the following equivalent definition of densely graded algebra.

Proposition 4.4. *Let G be a group, and $A = \oplus_{g\in G} A_g$ a G-graded algebra. Assume that A_g is finitely generated as a left A_e-module for all $g \in G$. Then A is a densely graded algebra if and only if there is a finite subset $\Upsilon \subseteq G \times G$ (Υ could be empty) such that*

(i) *for all $(g, h) \in (G \times G) \setminus \Upsilon$, $A_gA_h = A_{gh}$;*
(ii) *for all $g, h \in G$, A_{gh}/A_gA_h is finite dimensional.*

Proof. The "if" part is trivial. We next prove the "only if" part. Assume $A_gA_h \neq A_{gh}$. By Lemma 4.3, $A_gA_{g^{-1}} \neq A_e$ and $A_{h^{-1}}A_h \neq A_e$. Hence there could be finitely many elements $(g, h) \in G \times G$ such that $A_gA_h \neq A_{gh}$ by the condition (i). For $g \in G$, we have an exact sequence of right A_e-modules $0 \longrightarrow A_gA_{g^{-1}} \longrightarrow A_e \longrightarrow A_e/A_gA_{g^{-1}} \longrightarrow 0$. For any $h \in G$, applying $- \otimes_{A_e} A_{gh}$ on the sequence, we obtain the following exact sequence

$$A_gA_{g^{-1}} \otimes_{A_e} A_{gh} \longrightarrow A_{gh} \longrightarrow A_e/A_gA_{g^{-1}} \otimes_{A_e} A_{gh} \longrightarrow 0,$$

from which we obtain an isomorphism $A_{gh}/(A_gA_{g^{-1}}A_{gh}) \cong A_e/A_gA_{g^{-1}} \otimes_{A_e} A_{gh}$. By the condition (ii), $A_e/A_gA_{g^{-1}}$ is finite dimensional. Since A_{gh} is finitely generated A_e-module, $A_e/A_gA_{g^{-1}} \otimes_{A_e} A_{gh}$ is finite dimensional. Hence $A_{gh}/(A_gA_{g^{-1}}A_{gh})$ is finite dimensional. Note that $A_gA_{g^{-1}}A_{gh} \subseteq A_gA_h$. We finally obtain that A_{gh}/A_gA_h is finite dimensional for all $g, h \in G$. Hence A is a densely G-graded algebra. □

Let $A = \oplus_{g\in G}A_g$ be a G-graded algebra. Define a comodule action $\rho : A \to A \otimes \Bbbk G$ by $\rho(a) = a \otimes g$ for all $a \in A_g$ and $g \in G$. Then A is a right $\Bbbk G$-comodule algebra. On the contrary, any right $\Bbbk G$-comodule algebra A can be viewed as a G-graded algebra by setting components $A_g = \{a \in A | \rho(a) = a \otimes g\}$. The following result is a direct consequence of the proposition above.

Corollary 4.5. *With the hypotheses on A and G as in Proposition 4.4, A is a densely G-graded algebra if and only if A/A_e is a right $\Bbbk G$-dense Galois extension.*

Let $A = \oplus_{g\in G}A_g$ be a G-graded algebra. We denote by $\mathrm{GrMod}_G\, A$ the category of G-graded right A-module. One of the most important results for strongly graded algebras is Dade's Theorem (cf. [Da, Theorem 2.8], see also [NV]), which establishes an equivalence between the abelian categories $\mathrm{GrMod}_G\, A$ and $\mathrm{Mod}\, A_e$. In our case, these two categories are no longer equivalent in general. We next prove a weaker version of Dade's Theorem.

We may define two torsion classes in the abelian category $\mathrm{GrMod}_G\, A$. Let M be a G-graded right A-module. We call M a *G-graded torsion module* if the following condition is satisfied: for any cyclic graded submodule $N = mA$ generated by a homogeneous element m, we have that N_e is finite dimensional. Let $\mathcal{T}_G$ be the full subcategory of $\mathrm{GrMod}_G\, A$ consisting of all the graded torsion modules. For any G-graded right A-module N, we write $T_G(N)$ for the sum of all the torsion submodule of N. Then $T_G(N)$ is the largest torsion submodule of N. In this way, we indeed obtain a functor

$$T_G : \mathrm{GrMod}_G\, A \longrightarrow \mathcal{T}_G. \tag{4}$$

Another torsion class in $\mathrm{GrMod}_G\, A$ is the graded version of the one defined in previous section. Denote by $\mathrm{GrTor}_G\, A$ the full subcategory of $\mathrm{GrMod}_G\, A$ consisting of G-graded A-modules M such that mA is finite dimensional for each homogeneous element $m \in M$. One sees that $\mathrm{GrTor}_G\, A$ is a full subcategory of $\mathcal{T}_G$. In general, $\mathcal{T}_G$ is not equal to $\mathrm{GrTor}_G\, A$. However, we have the following result.

Lemma 4.6. *Let G be a finite group and A be a G-graded algebra which is also noetherian. If A is densely G-graded, then $\mathcal{T}_G = \mathrm{GrTor}_G\, A$.*

Proof. For $M \in \mathcal{T}_G$, let $m \in M$ be a homogenous element. Set $N = mA$. Then N is also a G-graded torsion module. Hence N_e is finite dimensional. Since A is noetherian and G is a finite group, A_e is a noetherian subalgebra of A and A_g is a finitely generated A_e-module on both sides for all $g \in G$. Then $N_e \otimes_{A_e} A_g$ is finite dimensional for all $g \in G$. The right multiplication of A on N gives a surjective map $N_e \otimes_{A_e} A_g \longrightarrow N_eA_g$. Hence N_eA_g is finite dimensional. For any $g \in G$, there is an exact sequence

$$0 \longrightarrow A_{g^{-1}}A_g \longrightarrow A_e \longrightarrow A_e/A_{g^{-1}}A_g \longrightarrow 0.$$

Applying the functor $N_g \otimes_{A_e} -$ to the exact sequence, we obtain the following exact sequence

$$N_g \otimes_{A_e} A_{g^{-1}}A_g \longrightarrow N_g \longrightarrow N_g \otimes_{A_e} (A/A_{g^{-1}}A_g) \longrightarrow 0.$$

Hence $N_g \otimes_{A_e} A_e/A_{g^{-1}}A_g \cong N_g/N_gA_{g^{-1}}A_g$. Since A is a densely G-graded algebra, $A_e/A_{g^{-1}}A_g$ is finite dimensional. Since A_g is finitely generated as a right A_e-module for all $g \in G$, N_g is a finitely generated A_e-module. Hence $N_g \otimes_{A_e} (A/A_{g^{-1}}A_g)$ is finite dimensional, which in turn implies that $N_g/N_gA_{g^{-1}}A_g$ is finite dimensional. Note that $N_gA_{g^{-1}}A_g \subseteq N_eA_g$. Hence N_g/N_eA_g is finite dimensional. As we already showed that N_eA_g is finite dimensional, it follows that N_g is finite dimensional for all $g \in G$. Hence $N = mA$ is finite dimensional as G is a finite group. Therefore, M is an object in $\mathrm{GrTor}_G A$. □

If A is a noetherian algebra, then both $\mathcal{T}_C$ and $\mathrm{GrTor}_G A$ are localizing Serre subcategories of $\mathrm{GrMod}_G A$. Hence we have quotient categories

$$\mathrm{GrMod}_G A/\mathcal{T}_G, \qquad \text{and} \qquad \mathrm{QGrMod}_G A := \frac{\mathrm{GrMod}_G A}{\mathrm{GrTor}_G A}.$$

The functor $(-)_e : \mathrm{GrMod}_G A \longrightarrow \mathrm{Mod}\, A_e$ sending $M = \oplus_{g\in G} M_g$ to M_e is an exact functor. One sees that $(-)_e$ sends objects in $\mathcal{T}_G$ to the objects in $\mathrm{Tor}\, A_e$. Hence it induces functors (we use the same symbol)

$$(-)_e : \mathrm{QGrMod}_G A \longrightarrow \mathrm{QMod}\, A_e, \qquad \text{and} \qquad (-)_e : \mathrm{GrMod}_G A/\mathcal{T}_G \longrightarrow \mathrm{QMod}\, A_e,$$

which make the following diagram commute

(5)
$$\begin{array}{ccc} \mathrm{QGrMod}_G A & \xrightarrow{(-)_e} & \mathrm{QMod}\, A_e, \\ \downarrow{\scriptstyle \pi} & \nearrow{\scriptstyle (-)_e} & \\ \mathrm{GrMod}_G A/\mathcal{T}_G & & \end{array}$$

where π is the natural projection functor.

Theorem 4.7 (Dade's Theorem)**.** *Let G be a finite group, and let $A = \oplus_{g\in G} A_g$ be a G-graded algebra. Assume that A is a noetherian algebra. Then the following are equivalent.*

(i) *A is a densely G-graded algebra.*
(ii) *For any finitely generated G-graded A-module $M = \oplus_{g\in G} M_g$, if M_e is finite dimensional, then M itself is finite dimensional.*
(iii) *The functor $(-)_e : \mathrm{QGrMod}_G A \longrightarrow \mathrm{QMod}\, A_e$ is an equivalence of abelian categories.*

Proof. Let $\Bbbk G$ be the group algebra and $H := \Bbbk G^*$ be its dual Hopf algebra. Then A is an H-module algebra, and $A^H = A_e$. Set $B = A\#H$. It is well known that any G-graded A-module M can be viewed as a right B-module, and vice versa. Hence we may (and we do) identify the abelian categories $\mathrm{Mod}\, B$ and $\mathrm{GrMod}_G A$. Under this point of view, the torsion subcategory $\mathrm{Tor}\, B$ is identified to $\mathcal{T}_G$, and the torsion functors $T : \mathrm{Mod}\, B \longrightarrow \mathrm{Tor}\, B$ and $T_G : \mathrm{GrMod}_G A \longrightarrow \mathcal{T}_G$ coincides. Hence we have $\mathrm{QGrMod}_G A = \mathrm{QMod}\, B$, and therefore, the functor $- \otimes_{\mathcal{B}} \mathcal{A} : \mathrm{QMod}\, B \longrightarrow \mathrm{QMod}\, A_e$ as defined by (2) is equivalent to $(-)_e : \mathrm{QGrMod}_G A \longrightarrow \mathrm{QMod}\, A_e$.

That (i) implies (iii) follows from Lemma 4.6, Corollary 4.5 and Theorem 2.3.

(iii) $\Longrightarrow$ (ii). By the commutative diagram (5) and Theorem 2.3, we see that the projection functor $\pi : \mathrm{QGrMod}_G A \longrightarrow \mathrm{GrMod}_G A/\mathcal{T}_G$ is an equivalence. Hence $\mathcal{T}_G = \mathrm{GrTor}_G A$. Let $M = \oplus_{g\in G} M_g$

be a finitely generated G-graded A-module. If M_e is finite dimensional, then $M \in \mathcal{T}_G = \mathrm{GrTor}_G A$. Since M is finitely generated, it follows that M is finite dimensional.

(ii) $\Longrightarrow$ (i). Let M be a finitely generated G-graded A-module. Then $T_G(M)$ is also a finitely generated G-graded A-module. Hence $T_G(M)_e$ is finite dimensional since $T_G(M)$ is a G-graded torsion module. By (ii) $T_G(M)$ is finite dimensional. Hence (i) follows from 4.5 and Theorem 2.3. □

5. Application I: Projective schemes associated to Veroness subalgebras

Throughout this section, $A = A_0 \oplus A_1 \oplus \cdots$ is an $\mathbb{N}$-graded algebra. We assume that A is noetherian and that A is locally finite, that is, $\dim_{\Bbbk} A_i < \infty$ for all $i \geq 0$.

Given an integer $d \geq 1$, the *dth Veroness subalgebra* is define to be the $\mathbb{N}$-graded algebra $A^{(d)}$ whose nth component is A_{nd}. Since A is noetherian, then $A^{(d)}$ is also noetherian. Similar, if M is a $\mathbb{Z}$-graded right A-module, we have $M^{(d)} = \oplus_{n\in\mathbb{Z}} M_{nd}$, which is a graded right $A^{(d)}$-module.

Let $\mathrm{GrMod}_{\mathbb{Z}} A$ be the category of the $\mathbb{Z}$-graded right A-modules. An object $M \in \mathrm{GrMod}_{\mathbb{Z}} A$ is said to be an $\mathbb{Z}$-graded torsion module if mA is finite dimensional for every $m \in M$. Let $\mathrm{GrTor}_{\mathbb{Z}} A$ to be the full subcategory of $\mathrm{GrMod}_{\mathbb{Z}} A$ consisting of all the $\mathbb{Z}$-graded torsion modules. Let $\mathrm{grmod}_{\mathbb{Z}} A$ to be the category of the finitely generated $\mathbb{Z}$-graded right A-modules, and let $\mathrm{grtor}_{\mathbb{Z}} A = \mathrm{grmod}_{\mathbb{Z}} A \bigcap \mathrm{GrTor}_{\mathbb{Z}} A$.

Since A is noetherian, $\mathrm{GrTor}_{\mathbb{Z}} A$ (resp. $\mathrm{grtor}_{\mathbb{Z}} A$) is a Serre subcategory of $\mathrm{GrMod}_{\mathbb{Z}} A$ (resp. $\mathrm{grmod}_{\mathbb{Z}} A$). Hence we have quotient category

$$\mathrm{Tail}\, A := \frac{\mathrm{GrMod}_{\mathbb{Z}} A}{\mathrm{GrTor}_{\mathbb{Z}} A} \qquad (\text{ resp. } \mathrm{tail}\, A := \frac{\mathrm{grmod}_{\mathbb{Z}} A}{\mathrm{grtor}_{\mathbb{Z}} A}),$$

which is usually called *the noncommutative projective scheme* associated to A (cf. [AZ, Ve]).

There are strong relations between the noncommutative projective scheme associated to A and that associated to $A^{(d)}$. Indeed, there is a natural functor

$$(-)^{(d)} : \mathrm{GrMod}_{\mathbb{Z}} A \longrightarrow \mathrm{GrMod}_{\mathbb{Z}} A^{(d)}, \quad M \mapsto M^{(d)}.$$

Restrictions to the finitely generated modules, we have

$$(-)^{(d)} : \mathrm{grmod}_{\mathbb{Z}} A \longrightarrow \mathrm{grmod}_{\mathbb{Z}} A^{(d)}.$$

Since $(-)^{(d)}$ is exact and sends torsion modules to torsion module, we obtain functors (use the same symbol)

$$(-)^{(d)} : \mathrm{Tail}\, A \longrightarrow \mathrm{Tail}\, A^{(d)}, \quad \text{and} \quad (-)^{(d)} : \mathrm{tail}\, A \longrightarrow \mathrm{tail}\, A^{(d)}.$$

The following result is fundamental in noncommutative geometry, which firstly appeared in [Ve] (see also, [AZ, Po, Mo]).

Theorem 5.1. *Assume A is generated by A_0 and A_1. Then $(-)^{(d)} : \mathrm{Tail}\, A \longrightarrow \mathrm{Tail}\, A^{(d)}$ and $(-)^{(d)} : \mathrm{tail}\, A \longrightarrow \mathrm{tail}\, A^{(d)}$ are equivalences of abelian categories.*

We will recover this result in the framework of the dense Galois extensions. Moreover, we will give a necessary and sufficient condition for the functor $(-)^{(d)} : \mathrm{Tail}\, A \longrightarrow \mathrm{Tail}\, A^{(d)}$ to be an equivalence.

Let d be a positive integer. For any integer $n \in \mathbb{Z}$, we have a unique presentation $n = qd + r$ such that $0 \leq r < d$. We define maps $\sigma : \mathbb{Z} \to \mathbb{Z}, n \mapsto q$ and $\delta : \mathbb{Z} \to \mathbb{Z}, n \mapsto r$.

Given a positive integer, let $\mathbb{Z}_d$ be the cyclic group $\mathbb{Z}/d\mathbb{Z} = \{\overline{0}, \overline{1}, \ldots, \overline{d-1}\}$. Set $\Gamma = \mathbb{Z} \times \mathbb{Z}_d$. Define a multiplication on Γ as follows:

$$(i, \overline{s}) + (j, \overline{t}) = (i + j + \sigma(s + t), \overline{\delta(s + t)}), \text{ where } 0 \leq s, t \leq d - 1.$$

Lemma 5.2. *With notation as above.*

(i) Γ, *together the multiplication defined above, is an abelian group.*
(ii) *Define a map* $\varphi : \Gamma \to \mathbb{Z}, (j, \overline{t}) \mapsto jd + t$, *where* $0 \leq t \leq d - 1$. *Then* φ *is an isomorphism of abelian groups.*

Proof. A straightforward check. □

Go back to the locally finite noetherian algebra $A = A_0 \oplus A_1 \oplus \cdots$. We may regraded A as follows so that A becomes a Γ-graded algebra. For $0 \leq t \leq d - 1$ and $j \geq 0$, set

$$D_{(j,\overline{t})} = A_{jd+t},$$

and set $D_{(j,\overline{t})} = 0$ for $j < 0$.

Proposition 5.3. *With notation as above.*

(i) $D = \oplus_{(j,\overline{t})\in\Gamma} D_{(j,\overline{t})}$, *together with the same multiplication as* A, *is a* Γ*-graded algebra.*
(ii) *If we set* $D_{\overline{s}} = \oplus_{i\in\mathbb{Z}} D_{(i,\overline{s})}$, *then* $D = \oplus_{\overline{s}\in\mathbb{Z}_d} D_{\overline{s}}$ *is a* $\mathbb{Z}_d$*-graded algebra.*

Proof. A straightforward check. □

Similar to the statement (ii) of the proposition above, given a Γ-graded right D-module M, if we set $M_{\overline{s}} = \oplus_{i\in\mathbb{Z}} M_{(i,\overline{s})}$, then $M = \oplus_{\overline{s}\in\mathbb{Z}_d} M_{\overline{s}}$ is a $\mathbb{Z}_d$-graded module. From this point of view, the category of Γ-graded right D-modules can be viewed as the category of graded $\mathbb{Z}_d$-graded modules with an additional grading. The following result is the translation of Dade's Theorem 4.7. Note that $D_{\overline{0}} = A^{(d)}$.

Proposition 5.4. *With notation as above, the following are equivalent:*

(i) $D = \oplus_{\overline{s}\in\mathbb{Z}_d} D_{\overline{s}}$ *is a densely* $\mathbb{Z}_d$*-graded algebra.*
(ii) *For any finitely generated* Γ*-graded* D*-module* M, *if* $M_{\overline{0}} = \oplus_{i\in\mathbb{Z}} M_{(i,\overline{0})}$ *is finite dimensional, then* M *is finite dimensional.*
(iii) $(-)_{\overline{0}} : \mathrm{QGrMod}_\Gamma D \longrightarrow \mathrm{Tail}\, A^{(d)}$ *is an equivalence of abelian categories.*

Proof. The same proof of Dade's Theorem 4.7 works well in this situation. □

We may explain the proposition above in a more detailed way. Define functors

$$\Phi : \mathrm{GrMod}_\mathbb{Z} A \longrightarrow \mathrm{GrMod}_\Gamma D$$

by setting $\Phi(M) = \bigoplus_{(j,\overline{t})\in\Gamma} M_{(j,\overline{t})}$ where $M_{(j,\overline{t})} = M_{jd+t}$, and

$$\Psi : \mathrm{GrMod}_\Gamma D \longrightarrow \mathrm{GrMod}_\mathbb{Z} A$$

by setting $\Psi(N) = \bigoplus_{n\in\mathbb{Z}} N_n$ where $N_n = N_{(\sigma(n),\overline{\delta(n)})}$.

Since the group Γ is isomorphic to $\mathbb{Z}$ by Lemma 5.2, one sees $\Phi \circ \Psi = \mathrm{id}$ and $\Psi \circ \Phi = \mathrm{id}$. Hence, we may (and we do) identify $\mathrm{GrMod}_{\mathbb{Z}} A$ and $\mathrm{GrMod}_{\Gamma} D$. By taking this point of view, we also have $\mathrm{grmod}_{\mathbb{Z}} A = \mathrm{grmod}_{\Gamma} D$, $\mathrm{Tail}\, A = \mathrm{QGrMod}_{\Gamma} D$ and $\mathrm{tail}\, A = \mathrm{qgrmod}_{\Gamma} D$, where $\mathrm{qgrmod}_{\Gamma} D = \frac{\mathrm{grmod}_{\Gamma} D}{\mathrm{GrTor}_{\Gamma} D \bigcap \mathrm{grmod}_{\Gamma} D}$.

Theorem 5.5. *Let $A = A_0 \oplus A_1 \oplus \cdots$ be a locally finite noetherian $\mathbb{N}$-graded algebra and let d be a fixed positive integer. The following are equivalent.*

(i) *There is an integer $p > 0$ such that, for all $n \geq p$ and $0 \leq s \leq d-1$,*

$$A_{nd} = \sum_{i+j=n-1} A_{id+s} A_{jd+d-s}.$$

(ii) *Given a finitely generated right graded A-module M, if $M^{(d)} = \oplus_{n\in\mathbb{Z}} M_{nd}$ is finite dimensional, then M itself is finite dimensional.*

(iii) $(-)^{(d)} : \mathrm{Tail}\, A \longrightarrow \mathrm{Tail}\, A^{(d)}$ *is an equivalence of abelian categories.*

(iv) $(-)^{(d)} : \mathrm{tail}\, A \longrightarrow \mathrm{tail}\, A^{(d)}$ *is an equivalence of abelian categories.*

Proof. By definition, the $\mathbb{Z}_d$-graded algebra D is densely graded if and only if $D_{\overline{0}}/D_{\overline{s}}D_{\overline{d-s}}$ is finite dimensional for all $\overline{s} \in \mathbb{Z}_d$, which is equivalent to the condition (i). Since A is noetherian, any finitely generated graded submodule of a right graded A-module M is still finitely generated. Equivalently, any finitely generated Γ-graded submodule of M when viewed as a Γ-graded D-module is finitely generated. Hence the condition (ii) in Proposition 5.4 is equivalent to the condition (ii) in this theorem. Since $\mathrm{Tail}\, A = \mathrm{QGrMod}_{\Gamma} D$, the functor $(-)^{(d)}$ is corresponding to the functor $(-)_{\overline{0}}$. Hence the condition (iii) in Proposition 5.4 is equivalent to the condition (iii) in this theorem. Hence (i) $\Longleftrightarrow$ (ii) $\Longleftrightarrow$ (iii) follows form Proposition 5.4.

Note that the functor $(-)^{(d)}$ sends object in $\mathrm{tail}\, A$ to the objects in $\mathrm{tail}\, A^{(d)}$. Below, we use π to denote the projection functors from $\mathrm{grmod}_{\mathbb{Z}} A$ (resp. $\mathrm{grmod}_{\mathbb{Z}} A^{(d)}$) to $\mathrm{tail}\, A$ (resp. $\mathrm{tail}\, A^{(d)}$). For any object $\mathcal{N}$ in $\mathrm{tail}\, A^{(d)}$, let $N \in \mathrm{grmod}_{\mathbb{Z}} A^{(d)}$ such that $\pi(N) = \mathcal{N}$. We view N as a $D_{\overline{0}}$-module. Let $M = N \otimes_{D_{\overline{0}}} D$. Then M is a Γ-graded finitely generated D-module. Since we identify $\mathrm{GrMod}_{\Gamma} D$ with $\mathrm{GrMod}_{\mathbb{Z}} A$, M is a finitely generated A-module. One sees that $M_{\overline{0}} = M^{(d)} = N$. Hence $(\pi(M))^{(d)} = \mathcal{N}$. Hence the condition (iii) implies the condition (iv).

(iv) $\Longrightarrow$ (ii). For any finitely graded right A-module M, write $\mathcal{M}$ for $\pi(M)$. Note that $\mathcal{M}^{(d)} = \pi(\oplus_{n\in\mathbb{Z}} M_{nd})$. If $\oplus_{n\in\mathbb{Z}} M_{nd}$ is finite dimensional, then $\mathcal{M}^{(d)} = \pi(\oplus_{n\in\mathbb{Z}} M_{nd}) = 0$. Since $(-)^{(d)}$ is an equivalence, we have $\mathcal{M} = 0$ in $\mathrm{tail}\, A$, and hence $M \in \mathrm{grtor}_{\mathbb{Z}} A$. Since M by assumption is finitely generated, M has to be finite dimensional. Therefore the condition (ii) holds. □

Remark 5.6. (i) If A is generated by A_0 and A_1, then one easily see that the condition (i) of Theorem 5.5 is satisfied. Hence in this case, we recover the classical Theorem 5.1.

(ii) More generally, if there is an integer $p > 0$ such that $A_i A_j = A_{i+j}$ for all $i + j \geq p$, then condition (i) of Theorem 5.5 is also satisfied. Hence in this case, $(-)^{(d)} : \mathrm{Tail}\, A \longrightarrow \mathrm{Tail}\, A^{(d)}$ is an equivalence of abelian categories.

(iii) Mori also provided some equivalent conditions for the functor $(-)^{(d)}$ to be an equivalence in [Mo, Theorem 3.5]. Note that in that paper, A is assumed to be a coherent connected graded algebra satisfying further homological conditions. In our case, we only assume that A is a locally

finite noetherian algebra. Moreover, we obtain the result under the framework of Hopf dense Galois extension, which is a new way to understand noncommutative projective schemes.

6. Application II: Noncommutative isolated singularities

Throughout this section, $A = \oplus_{n\in\mathbb{N}} A_n$ is always a locally finite noetherian $\mathbb{N}$-graded algebra. Following [Ue], A is called a *graded isolated singularity* if the abelian category $\operatorname{tail} A$ has finite global dimension, that is, there is an integer $p \geq 0$ such that $\operatorname{Ext}^i_{\operatorname{tail} A}(\mathcal{M},\mathcal{N}) = 0$ for all $i > p$ and all $\mathcal{M},\mathcal{N} \in \operatorname{tail} A$.

Let H be a finite dimensional semisimple Hopf algebra. Assume that A is a left H-module algebra, and that H-action preserves the grading of A. Then $B := A\#H$ is also an $\mathbb{N}$-graded algebra by setting the degree of elements of H to be zero. Since H is finite dimensional, A can be viewed as a right H^*-comodule algebra. Note that the comodule action $\rho : A \longrightarrow A \otimes H^*$ also preserves the grading of A. We may replace the module categories in Theorem 2.3 by the categories of $\mathbb{Z}$-graded modules. Then Theorem 2.3 reads as follows.

Theorem 6.1. *Let $A = A_0 \oplus A_1 \oplus \cdots$ be a locally finite noetherian $\mathbb{N}$-graded algebra, and let H be a finite dimensional semisimple Hopf algebra. Assume that A is a left H-module and the H-action preserves the grading of A. Set $B := A\#H$ and $R := A^H$, and let t be the integer of H such that $\varepsilon(t) = 1$. Then the following are equivalent.*

(i) *A/R is a right H^*-dense Galois extension.*
(ii) *For any finitely generated $\mathbb{Z}$-graded right B-module, if Mt is finite dimensional, then M itself is finite dimensional.*
(iii) *B/AtA is finite dimensional.*
(iv) *$- \otimes_{\mathcal{B}} \mathcal{A} : \operatorname{Tail} B \longrightarrow \operatorname{Tail} R$ is an equivalence of abelian categories.*
(v) *$- \otimes_{\mathcal{B}} \mathcal{A} : \operatorname{tail} B \longrightarrow \operatorname{tail} R$ is an equivalence of abelian categories.*

Proof. The statements (i), (iii) and (iv) are exactly the graded versions of the statements (i), (iii) and (iv) of Theorem 2.3. The graded version of the statement (ii) of Theorem 2.3 can be stated as follows: for any finitely generated $\mathbb{Z}$-graded right B-module M, $T_{\mathbb{Z}}(M)$ is finite dimensional, where $T_{\mathbb{Z}}(M)$ is the largest $\mathbb{Z}$-graded B-submodule of M such that $T_{\mathbb{Z}}(M)t$ is finite dimensional. Since B is noetherian, this is equivalent to the statement (ii) is this theorem. Hence (i), (ii), (iii) and (iv) are equivalent.

That (iv) implies (v) is because $\operatorname{tail} A$ (resp. $\operatorname{tail} R$) is full subcategory of $\operatorname{Tail} A$ (resp. $\operatorname{Tail} R$), and the restriction of $- \otimes_{\mathcal{B}} \mathcal{A}$ to $\operatorname{tail} A$ is densely onto $\operatorname{tail} R$.

(v) $\Longrightarrow$ (ii). Let M be a finitely generated $\mathbb{Z}$-graded right B-module. If Mt is finite dimensional, then $\pi(M) \otimes_{\mathcal{B}} \mathcal{A} \cong \pi(Mt) = 0$ in $\operatorname{tail} R$. Since $- \otimes_{\mathcal{B}} \mathcal{A}$ is an equivalence, $\pi(M)$ is finite dimensional. Hence M is finite dimensional. □

With assumptions as in Theorem 6.1, we obtaing the following.

Corollary 6.2. *If A is of finite global dimension and A/A^H is a right H^*-dense Galois extension, then A^H is a graded isolated singularity.*

Proof. If A is of finite global dimension, then so is B. Then the global dimension of $\operatorname{tail} B$ is finite. By Theorem 6.1, $\operatorname{tail} A^H$ has finite global dimension, and hence A^H is a graded isolated singularity. □

Remark 6.3. (i) Note that the graded version of Theorem 3.8 also holds. We leave it to the reader to write it down.

(ii) A general theory of graded isolated singularities was established by Ueyama [Ue] and Mori-Ueyama [MU]. Many interesting examples were presented in those papers in the case that A is a Artin-Schelter regular algebra and H is a cyclic group algebra.

(iii) It is well known that finite subgroups of $SL_2(\Bbbk)$ acting on the polynomial algebra $\Bbbk[x, y]$ yield isolated singularities. As for the noncommutative case, finite dimensional Hopf actions on quantum planes are well understood by Chan, Kirkman, Walton and Zhang in [CKWZ1]. If G is a finite group which acts homogeneous on a quantum plane A and the homological determinant of G is trivial, then the condition (iii) of Theorem 6.1 is always satisfied (cf. [Ue]). Hence the invariant subalgebra A^G is graded isolated by Corollary 6.2.

Recall from [MM] that a locally finite noetherian $\mathbb{N}$-graded algebra A is called an *Artin-Schelter Gorenstein algebra* if $\operatorname{injdim} A_A = \operatorname{injdim}_A A = n < \infty$, and

$$\operatorname{Ext}^i_A(A_0, A) \cong \begin{cases} 0, & i \neq n \\ A_0^*(l), & i = n \end{cases}$$

both in $\operatorname{GrMod}_{\mathbb{Z}} A$ and in $\operatorname{GrMod}_{\mathbb{Z}} A^{op}$, where where (l) is the degree shift functor.

Corollary 6.4. *Let H be a semisimple Hopf algebra, and let A be an $\mathbb{N}$-graded left H-module algebra. Assume that the H-action preserves the grading of A, and that A is an Artin-Schelter Gorenstein algebra with injective dimension $\operatorname{injdim}_A A = \operatorname{injdim} A_A = n \geq 2$. If A/A^H is a H^*-dense Galois extension, then the natural map*

$$A\#H \longrightarrow \operatorname{End}(A_{A^H}), a\#h \mapsto [b \mapsto a(h \cdot b)]$$

is an isomorphism of $\mathbb{N}$-graded algebras.

Proof. Since A is Artin-Schelter Gorenstein and the injective dimension of A is not less than 2, it follows that $\operatorname{depth} A_A \geq 2$ (we remark that in graded case, the depth of A is considered in $\operatorname{GrMod}_{\mathbb{Z}} A$). By the graded version of Theorem 3.8 (cf. Remark 6.3 (i)), we have $A\#H \cong \operatorname{End}(A_{A^H})$ as graded algebra. □

Finally, we remark that a similar result was already obtained in [MU, Theorem 3.7] under the much stronger assumptions that A is a noetherian Artin-Schelter regular algebra and H is a finite group algebra.

Acknowledgement. The first author is supported by NSFC (No. 11571239) and NSF of Zhejiang Province (No. LY14A010006). The second and the third authors are supported by an FWO-grant.

References

[AZ] M. Artin, J. J. Zhang, Noncommutative projective schemes, Adv. Math. 109 (1994), 228–287.

[Da] E. C. Dade, Group graded rings and modules, Math. Z. 174 (1980), 241–262.

[KT] H. F. Kreimer, M. Takeuchi, Hopf algebras and Galois extensions of an algebra, Indiana Univ. Math. J. 30 (1981), 675–692.

[CFM] M. Cohen, D. Fischman, S. Montgomery, Hopf Galois extensions, Smash products, and Morita equivalence, J. Algebra 133 (1990), 335–372.

[CKWZ1] K. Chan, E. Kirkman, C. Walton, J. J. Zhang, Quantum binary polyhedral groups and their actions on quantum planes, J. Reine Angew. Math. (Crelles Journal), DOI: 10.1515/crelle-2014-0047, 2014.

[CKWZ2] K. Chan, E. Kirkman, C. Walton, J. J. Zhang, McKay Correspondence for semisimple Hopf actions on regular graded algebras, preprint, 2015.

[EW] P. Etingof, C. Walton, Semisimple Hopf actions on commutative domains, Adv. math. 251 (2014), 47–61.

[Ga] N. Gao, Hopf extensions of CM-finite Artin aglebras, Algebr. Reprent. Theory 16 (2013), 605–613.

[HVZ] J.-W. He, F. Van Oystaeyen, Y. Zhang, Hopf algebra actions on differential graded algebras and applications, Bull. Belg. Math. Soc. Simon Stevin 18 (2011), 99–111.

[JZ] P. Jorgensen, J. J. Zhang, Gourmets guide to Gorensteinness, Adv. Math. 151 (2000), 313–345.

[KKZ1] E. Kirkman, J. Kuzmanovich, J.J. Zhang, Gorenstein subrings of invariants under Hopf algebra actions, J. Algebra 322 (2009), 3640–3669.

[KKZ2] E. Kirkman, J. Kuzmanovich, J.J. Zhang, Shephard-Todd-Chevalley theorem for skew polynomial rings, Algebr. Represent. Theory 13 (2010), 127–158.

[KKZ3] E. Kirkman, J. Kuzmanovich, J. J. Zhang, Noncommutative complete intersections, J. Algebra 429 (2015), 253–286.

[MM] H. Minamoto, I. Mori, The structure of AS-Gorenstein algebras, Adv. Math. 226 (2011), 4061–4095.

[Mon] S. Montgomery, Hopf Algebras and Their Actions on Rings, CBMS Reg. Conf. Ser. Math. 82, Amer. Math. Soc., Providence, RI, 1993.

[Mo] I. Mori, McKay type correspondence for AS-regular algebras, J. London Math. Soc. (2) 88 (2013), 97–117.

[MU] I. Mori, K. Ueyama, Ample group action on AS-regular algebras and noncommutative graded isolated singularities, to appear at Trans. Amer. Math. Soc., arXiv:1404.5045.

[NV] C. Năstăsescu, F. Van Oystaeyen, Methods of Graded Rings, Lect. Notes Math. 1836, Springer, 2004.

[Po] A. Polishchuk, Noncommutative proj and coherent algebras, Math. Res. Lett. 12 (2005), 63–74.

[PP] N. Popescu, L. Popescu, Theory of Categories, Editura Academiei, Bucureşti, România and Sijthoff & Noordhoff International Publishers, 1979.

[St] B. Stenström, Rings and Modules of Quotients, Lect. Notes Math. 237, Springer-Verlag, 1971.

[Ue] K. Ueyama, Graded maximal Cohen-Macaulay modules over noncommutative graded Gorenstein isolated singularities, J. Algebra 383 (2013), 85–103.

[VZ] F. Van Oystaeyen, Y. Zhang, Introduction functors and stable Clifford theory for Hopf algebras, J. Pure Appl. Algebra 107 (1996), 337–351.

[Ve] A. B. Verevkin, On the noncommutative analogue of the category of coherent sheaves on a projective scheme, Amer. Math. Soc. Transl. (2) 151 (1992), 41–53.

[WW] C. Walton, X. Wang, On quantum groups associated to non-Noetherian regular algebras of dimension 2, arXiv:1503.09185.

J.-W. He

Department of Mathematics, Hangzhou Normal University, 16 Xuelin Rd, Hangzhou Zhejiang 310036, China

E-mail address: jwhe@hznu.edu.cn

F. Van Oystaeyen

Department of Mathematics and Computer Science, University of Antwerp, Middelheimlaan 1, B-2020 Antwerp, Belgium

E-mail address: fred.vanoystaeyen@ua.ac.be

Y. Zhang
Department WNI, University of Hasselt, Universitaire Campus, 3590 Diepenbeek, Belgium

E-mail address: yinhuo.zhang@uhasselt.be